



## INTRAUTERINE DEVICE (IUD) INSERTION CONSENT

I request a:  Mirena IUD  Liletta IUD  Kyleena IUD  Paragard IUD

I understand the following (please initial):

\_\_\_\_\_ I may have a pregnancy test before the IUD is inserted. If I had unprotected sex within the past 7 days, the pregnancy test may read negative when an early pregnancy is starting.

\_\_\_\_\_ The Mirena and Liletta protect against pregnancy for 7 years. The Kyleena protects against pregnancy for 5 years. The Paragard protects against pregnancy for 10 years.

\_\_\_\_\_ The possible risks of IUD placement include infection, bleeding, allergic reaction, perforation of (poking a hole in) the uterus, expulsion (falling out) or incorrect placement (placed in the uterine muscle or abdominal cavity) of the IUD. Perforation or incorrect placement will most likely require removal of the IUD including possible surgery if the IUD is in the abdominal cavity. These risks are higher in women who have recently finished a pregnancy or who are currently breastfeeding.

\_\_\_\_\_ I may have irregular bleeding and cramping for the first 3 months after the IUD is inserted. Ibuprofen, naproxen, or a heating pad may help with these symptoms.

\_\_\_\_\_ The IUD does not protect against sexually transmitted infections (STIs). I should use condoms to protect myself against STIs.

\_\_\_\_\_ With the Mirena, Kyleena, and Liletta IUDs, my periods may get lighter or go away. This is common and not dangerous. With the Paragard IUD, my periods may get heavier or last longer.

\_\_\_\_\_ I understand the risks, possible side effects, and know what to expect after the IUD is inserted.

\_\_\_\_\_

Patient Name

\_\_\_\_\_/\_\_\_\_\_/\_\_\_\_\_

Date

\_\_\_\_\_

Patient/Health Care Agent/Guardian/Relative Signature

\_\_\_\_\_

Physician Signature